

Flexor tendon repair zones 1-3 Modified Duran (Passive)

This protocol is intended to provide the clinician with a guideline for the postoperative rehabilitation course of a patient who has undergone flexor digitorum superficialis and/or flexor digitorum profundus repairs in Zones 1-3 and whose surgeon has specifically ordered a Modified Duran or passive protocol. General time frames are given for reference to the average, but individual patients will progress at different rates depending on their age, comorbidities, pre-surgical range of motion, strength, health/functional status, rehabilitation compliance, learning barriers and complications. Specific time frames, restrictions and precautions are given to protect healing tissues and surgical reconstruction.

After 4 weeks, use Strickland's Percentage to track progress. (.stricklands smartphrase)

((Active PIP + DIP) - PIP extension lag) divided by 175) x 100 = % of normal active PIP and DIP motion

- Excellent = 85-100%
- Good = 70-84%
- Fair = 50-69%
- Poor = <50%

*Measurements were taken actively, not after place and hold

AROM	Date
PIP	***/**
DIP	***/**
TAM	***
STRICKLAND'S %	***%
STRICKLAND'S RATING	***

Postoperative Guidelines

First post-operative visit should be 3-7 days post operatively.

If patient is diabetic, a smoker or has other underlying medical conditions to have slow healing, they may need extended orthotic time to achieve a satisfactory outcome.

For Zone 1 repairs, in addition to Wrist Hand Finger Orthosis (WHFO), consider dorsal blocking splint positioning DIP of affected digit in 45 degrees flexion

Return to Work

Patients who work as manual laborers are normally able to return to work approximately 3-6 months after surgery. Those who have sedentary professions are normally able to return to work 6-8 weeks after surgery.

The following should be taken into consideration:

- Extent of Injuries
- Type of work
- Surgeon's approval
- Postoperative complications

Phase I (surgery to 3-5 days after surgery)

Rehabilitation appointments	<ul style="list-style-type: none"> • 2-4 times for the first month
Rehabilitation goals and priorities	<ul style="list-style-type: none"> • Protect tendon repair • Teach PROM • Manage edema • Wound healing • One-handed activities of daily living (ADLs)
Orthosis	<ul style="list-style-type: none"> • Custom thermoplastic dorsal blocking WHFO • Wrist in neutral to slight extension, MCPs 70 degrees flexion, IPs full extension • If needed, apply light compression (example: coban or compressogrip) for edematous finger(s). • For Dr. Salyapongse patients: no strap blocking IP flexion
Suggested therapeutic exercises	<p>Performed within splint at home, 10 reps each every 1-2 hours (no less than 5x/day)</p> <ul style="list-style-type: none"> ○ Passive flexion and active/passive extension. ○ Passive composite fist ○ Passive hook fist.
Precautions	<ul style="list-style-type: none"> • Orthosis on 24/7 (including bathing), remove ONLY for careful hygiene at sink
Progression criteria	<ul style="list-style-type: none"> • Full PROM of finger(s): passive flexion pulp to palm

Flexor Tendon Repairs Modified Duran Zone 1-3

Phase II (4-6 weeks post-op)

Rehabilitation appointments	<ul style="list-style-type: none"> ○ Twice per week
Rehabilitation goals and priorities	<ul style="list-style-type: none"> ● Tendon gliding to optimize intrinsic tendon healing and minimize extrinsic scarring ● Manage edema ● Manage scar
Orthosis	<ul style="list-style-type: none"> ● Allow removal of orthosis to shower, emphasizing not to use affected hand while bathing ● Orthosis on 24/7, remove ONLY for careful hygiene ● Modify orthosis to wrist at 30 degrees extension, MCPs at 30 degrees flexion, and IPs in extension.
Suggested therapeutic exercises	<ul style="list-style-type: none"> ● Measure using Strickland’s to determine progression of exercises according to flexor tendon pyramid ● Passive flexion and extension warm-up-FIRST ● Gentle active tendon gliding exercises, focusing on hook, composite and straight fist. ● Block MCP in full extension and perform full active IP extension ● Synergistic wrist motion: gravity-assisted wrist flexion (pronated) followed by active wrist extension to splint
Precautions	<ul style="list-style-type: none"> ○ No functional use of hand. ○ No composite finger/wrist extension

Flexor tendon repairs Modified Duran Zone 1-3

Phase III (6-8 weeks post-op)

Rehabilitation appointments	<ul style="list-style-type: none"> Twice per week
Rehabilitation goals and priorities	<ul style="list-style-type: none"> Progress to full AROM Being using hand for specific light ADL tasks while seated
Orthosis	<ul style="list-style-type: none"> Discontinue splint
Suggested therapeutic exercises	<p>Continue progression using Strickland’s measurement and clinical reasoning, based on individual patient response</p> <p>Blocking: stabilize finger on lateral surfaces (to minimize work of flexion)</p> <ul style="list-style-type: none"> Begin with blocking MPJ in slight flexion Progress to blocking MPJ in full extension Progress to blocking PIPJ in slight flexion Progress to blocking PIPJ in full extension
Precautions	No lifting/pulling/pushing/gripping
Progression criteria	<p>Strickland's Percentage= ((Active PIP + DIP flexion) - (PIP + DIP extension lag)) divided by 175) x 100 = % of normal active PIP and DIP motion</p> <ul style="list-style-type: none"> Excellent = 85-100% Good = 70-84% Fair = 50-69% Poor = <50% <p>*Measurements taken actively, not after place and hold</p>

Flexor tendon repairs Modified Duran zone 1-3

Phase IV (8-12 weeks post-op)

Rehabilitation appointments	<ul style="list-style-type: none"> • Depending on scar, frequency varies from 2 times per week to 2 times per month
Rehabilitation goals and priorities	<ul style="list-style-type: none"> • Progress to using hand for light ADL/IADL tasks • Full AROM • Minimize scarring • Gradually increase strength
Suggested therapeutic exercises	<ul style="list-style-type: none"> • Passive composite finger/wrist extension <ul style="list-style-type: none"> ○ Consider spring extension splint or thermoplastic night splint as needed if flexion contracture at PIPJ • Consider otoform mold or silicone gel sheeting for scar • Isolated IP flexion activities <ul style="list-style-type: none"> ○ Scrunching washcloth on table with palm flat ○ Picking up progressively smaller objects between pulp of affected finger(s) and palm ○ Rotating cylindrical objects on tabletop between thumb and affected finger(s) ○ Actively moving between hook fist and composite fist while holding a pen or highlighter • Initiate strengthening if needed (light therapeutty)
Precautions	<ul style="list-style-type: none"> • No dynamometer testing • No torque/resistance activity allowed

References

- Cannon, N. M., Beal, B., & Walters, K. (2001). Diagnosis and treatment manual for physicians and therapists. *Indianapolis: The Hand Center of Indiana PC.*
- Chesney, A., Chauhan, A., Kattan, A., Farrokhyar, F., & Thoma, A. (2011). Systematic review of flexor tendon rehabilitation protocols in zone II of the hand. *Plastic and Reconstructive Surgery, 127*(4), 1583-1592.
- Clancy, S. P., & Mass, D. P. (2013). Current flexor and extensor tendon motion regimens. *Hand Clinics, 29*(2), 295-309.
- Coates II, R. W., Echevarria-Ore, J. C., & Mass, D. P. (2005). Acute flexor tendon repairs in zone II. *Hand Clinic, 21*, 173-179. Doi: 10.1016/j.hcl.2004.11.001
- Edsfeldt, S., Rempel, D., Kursa, K., Diao, E., & Lattanza, L. (2015). In vivo flexor tendon forces generated during different rehabilitation exercises. *Journal of Hand Surgery European Volume, 40*(7), 705-710. Doi: 10.1177/1753193415591491
- Frueh, F. S., Kunz, V. S., Gravestock, I. J., Held, L., Haefeli, M., Giovanoli, P., & Calcagni, M. (2014). Primary flexor tendon repair in zones 1 and 2: early passive mobilization versus controlled active motion. *The Journal of Hand Surgery, 39*(7), 1344-1350.
- Higgins, A., Lalonde, D. H., Bell, M., Mckee, D., & Lalonde, J. F. (2010). Avoiding flexor tendon repair rupture with intraoperative total active movement examination. *Plastic Reconstructive Surgery, 126*(3), 941-945. Doi: 10.1097/PRS.0b013e3181e60489
- Higgins, A., & Lalonde, D. H. (2016). Flexor Tendon Repair Postoperative Rehabilitation: The Saint John Protocol. *Plastic and Reconstructive Surgery Global Open, 4*(11).
- Howell, J. W., & Peck, F. (2013). Rehabilitation of flexor and extensor tendon injuries in the hand: current updates. *Injury, 44*(3), 397-402.
- Lalonde, D. (2013). How the wide awake approach is changing hand surgery and hand therapy: Inaugural AAHS sponsored lecture at the ASHT meeting, San Diego, 2012. *Journal of Hand Therapy, 26*, 175-178. Doi: 10.1016/j.jht.2012.12.002
- Neiduski, R. (2016). Motion and measurement: Achieving optimal outcomes after flexor tendon repair. *UW-Milwaukee Continuing Education Course #9025.*
- Orkar, K. S., Watts, C., & Iwuagwu, F. C. (2011). A comparative analysis of the outcome of flexor tendon repair in the index and little fingers: Does the little finger fare worse? *Journal of Hand Surgery European Volume, 37*(1), 20-26. Doi: 10.1177/1753193411413661
- Tang, J. B. (2007). Indications, methods, postoperative motion and outcome evaluation of primary flexor tendon repairs in zone 2. *Journal of Hand Surgery European Volume, 32*(2), 118-129. Doi: 10.1016/j.jhsb.2006.12.009

- Skirven, T. M., Osterman, A. L., Fedorczyk, J., & Amadio, P. C., Feldscher, S.B., Shin, E.K. (2021). *Rehabilitation of the Hand and Upper Extremity 7th edition, 2-Volume Set E-Book: Expert Consult*. Elsevier Health Sciences.
- Skirven, T. M., Osterman, A. L., Fedorczyk, J., & Amadio, P. C. (2011). *Rehabilitation of the Hand and Upper Extremity 6th edition, 2-Volume Set E-Book: Expert Consult*. Elsevier Health Sciences.
- Starr, H. M., Snoddy, M., Hammond, K. E., & Seiler, J. G. (2013). Flexor tendon repair rehabilitation protocols: a systematic review. *The Journal of Hand Surgery*, 38(9), 1712-1717.
- Tang, J. B., Amadio, P. C., Guimberteau, J. C., & Chang, J. (2012). *Tendon Surgery of the Hand E-Book*. Elsevier Health Sciences.
- Tang, J. B., Chang, J., Elliot, D., Lalonde, D. H., Sandow, M., & Vögelin, E. (2014). IFSSH flexor tendon committee report 2014: from the IFSSH flexor tendon committee (chairman: Jin Bo Tang). *Journal of Hand Surgery (European Volume)*, 39(1), 107-115.
- Wu, Y. F., & Tang, J. B. (2014). Recent developments in flexor tendon repair techniques and factors influencing strength of the tendon repair. *Journal of Hand Surgery (European Volume)*, 39(1), 6

These rehabilitation guidelines were developed collaboratively between UW Health and UnityPoint Health - Meriter Rehabilitation and the UW Health Orthopedic Surgeons.

Content is for informational purposes only and does not replace the guidance, diagnostic or treatment options or educational materials your healthcare provider gives you. Call your health provider immediately if you think you may have a medical emergency. Always seek the advice of your health provider prior to starting any new treatment and contact them immediately with any medical emergency.